

Improving Survivorship for Cancer Patients

Latest Research Shows Lymphedema is a Risk of Cancer Treatment



million
1.8

newly diagnosed cancer patients every year¹


■ 50% of cancer patients are at risk for limb lymphedema¹⁻³



1 in 3

at risk cancer patients will develop lymphedema^{2,3}, e.g.

- Breast cancer
- Melanoma
- Pelvic area cancer



75%

of lymphedema is diagnosed within three years of treatment⁴:

- Lymph node surgery
- Nodal irradiation
- Taxane-based chemotherapy

Early Detection & Intervention are Key to Preventing Lymphedema

95% ↓

of patients did not progress to chronic lymphedema with early detection using L-Dex® and intervention⁵

The opportunity to prevent lymphedema is before noticeable swelling exists

Testing for Early Lymphedema is Easy

L-Dex® testing with the SOZO® Digital Health Platform



Learn more
www.impedimed.com/improving-survivorship

↓
**Fast
Non-invasive
Easy-to-use**
↑



REFERENCES

1) American Cancer Society's Cancer Facts and Figures, 2019. <https://www.cancer.org/content/dam/cancer-org/research/cancer-facts-and-statistics/annual-cancer-facts-and-figures/2019/cancer-facts-and-figures-2019.pdf>. 2) Shaitelman SF, et al. Recent Progress in the Treatment and Prevention of Cancer-Related Lymphedema. CA Cancer J Clin. 2015;00:00-00. 3) Brown JC, et al. The Prevalence of Lymphedema Symptoms among Long-Term Cancer Survivors with or at-risk for Lower Limb Lymphedema. Am J Phys Med Rehabil. 2013 March; 92(3): 223-231. doi:10.1097/PHM.0b013e31826edd97. 4) Herd-Smith, A., et al., Prognostic factors for lymphedema after primary treatment of breast carcinoma. Cancer, 2001. 92(7): p. 1783-7. 5) Ridner SH, et al. A Randomized Trial Evaluating Bioimpedance Spectroscopy Versus Tape Measurement for the Prevention of Lymphedema Following Treatment for Breast Cancer: Interim Analysis. Ann Surg Oncol 2019; <https://doi.org/10.1245/s10434-019-07344-5>.